



UNIVERSITI PUTRA MALAYSIA

**STUDIES IN KEDAH-KELANTAN, BRAHMAN AND
CROSSBRED BULLS AT PUBERTY AND SEXUAL MATURITY**

ISMAYA

FPV 1987 1

STUDIES IN KEDAH–KELANTAN, BRAHMAN AND
CROSSBRED BULLS AT PUBERTY AND SEXUAL MATURITY

By

ISMAYA

A thesis submitted in partial fulfilment of the requirements
for the degree of Master of Science in the
Faculty of Veterinary Medicine and Animal Science
Universiti Pertanian Malaysia




It is hereby certified that we have read this thesis entitled 'Studies in Kedah-Kelantan, Brahman and Crossbred Bulls at Puberty and Sexual Maturity' by Ismaya, and in our opinion it is satisfactory in terms of scope, quality, and presentation as partial fulfilment of the requirements for the degree of Master of Science


.....
ALANG P. ZAINUDDIN, Ph.D.

Assoc. Professor/Dean of Graduate Studies
Universiti Pertanian Malaysia
(Chairman Board of Examiners)


.....
K.W. ENTWISTLE, Ph. D.

Professor
Graduate School of Tropical Veterinary Science
James Cook University of North Queensland
Townsville, AUSTRALIA
(External Examiner)


.....
ADNAN BIN SULONG, Ph.D.

Lecturer
Faculty of Veterinary Medicine and Animal Science
Universiti Pertanian Malaysia
(Internal Examiner)


.....
TAN HOCK SENG, Ph.D.

Associate Professor
Faculty of Veterinary Medicine and Animal Science
Universiti Pertanian Malaysia
(Internal Examiner/Supervisor)

This thesis was submitted to the Senate of Universiti
Pertanian Malaysia and was accepted as partial fulfilment of
the requirements for the degree of Master of Science.

Date: **15 JUL 1987**


Dean of Graduate Studies

Dedicated to Tini, H.S.



ACKNOWLEDGEMENT

Praise be to the Almighty Allah. Had it not been His will, the completion of this study would not have been possible.

I would like to take this opportunity to express my deep gratitude and sincere appreciation to Associate Professor Dr. Tan Hock Seng, my supervisor, for his invaluable guidance, continuous supervision, constructive criticism, encouragement, understanding and helpful advice throughout the course of this study.

My gratitude and appreciation are also extended to Associate Professor Dr. Tuan Arifteen Bongso, my co-supervisor, for his advice.

I am grateful to the Dean of Faculty of Veterinary Medicine and Animal Science for all facilities provided during the course of this work. I am also grateful to the Malaysian Technical Cooperation Programme (MTCP) for financial support.

Sincere thanks are due to Dr. A. Rahim and staff of Darabif Sdn. Bhd., Pahang as well as Encik Tukiran and co-workers of Beef Unit, Universiti Pertanian Malaysia for their assistance. Thanks are also extended to Mrs. Zainab bte Nasri for her assistance and technical advice.



Sincere thanks are also due to the Government of the Republic of Indonesia and the Universitas Gadjah Mada, Yogyakarta, for granting the study leave.

Lastly, I would like to dedicate this thesis to my parents and Dikiyati Yansumayutipa. Their understanding and patience has enabled me to complete this work.

TABLE OF CONTENTS

| | Page |
|--|-------|
| ACKNOWLEDGEMENT | iii |
| TABLE OF CONTENTS | v |
| LIST OF TABLES | viii |
| LIST OF FIGURES | x |
| LIST OF APPENDICES | xii |
| LIST OF ABBREVIATIONS | xiii |
| ABSTRACT | xv |
| CHAPTER 1 INTRODUCTION | 1 |
| CHAPTER 2 REVIEW OF LITERATURE | 5 |
| Puberty | 5 |
| Assessment of Puberty | 7 |
| Body Weight | 7 |
| Heart Girth | 7 |
| Scrotal Circumference | 7 |
| Sperm Production | 11 |
| Libido | 13 |
| Testosterone Concentration | 14 |
| Factors Affecting Puberty | 18 |
| Effect of Breed | 18 |
| Effect of Age | 20 |
| Effect of Nutrition | 21 |



| | Page |
|---|------|
| Spermatogenesis | 23 |
| The Process of Spermatogenesis | 23 |
| Hormonal Control of Spermatogenesis | 24 |
| Spermatozoa | 26 |
| CHAPTER 3 MATERIALS AND METHODS | 28 |
| Experimental Animals | 28 |
| Measurement of Heart Girth, Body Weight, Scrotal Circumference, Testis Length, Testis Width and Testes Volume | 29 |
| Blood Sampling and GnRH Administration | 29 |
| Semen Collection | 30 |
| Semen Evaluation | 31 |
| Sperm Motility | 31 |
| Sperm Concentration | 31 |
| Percentage of Live Sperm | 32 |
| Radioimmunoassay (RIA) | 33 |
| Chemicals and Apparatus | 33 |
| Label and Antisera | 34 |
| Preparation of Reagents | 34 |
| Radioimmunoassay Procedure | 35 |
| Quality Control | 37 |
| Sensitivity | 38 |
| Statistical Analysis | 39 |



| | Page |
|--|------|
| CHAPTER 4 RESULTS | 40 |
| Heart Girth | 40 |
| Body Weight | 46 |
| Testis Size | 52 |
| Scrotal Circumference | 52 |
| Testes Volume | 59 |
| Testis Length | 62 |
| Testis Width | 69 |
| Sperm Production | 73 |
| Plasma Testosterone Levels in Kedah- Kelantan Bulls | 76 |
| CHAPTER 5 DISCUSSION | 89 |
| Heart Girth | 89 |
| Body Weight | 90 |
| Testis Size | 94 |
| Scrotal Circumference | 94 |
| Testes Volume | 98 |
| Testis Length | 99 |
| Testis Width | 100 |
| Sperm Production | 101 |
| Plasma Testosterone Levels in Kedah- Kelantan Bulls | 103 |
| CHAPTER 6 CONCLUSION | 107 |
| BIBLIOGRAPHY | 110 |
| APPENDICES | 119 |



LIST OF TABLES

| Table | | Page |
|-------|--|------|
| 1 | Comparison of heart girth in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 41 |
| 2 | Correlation and regression between heart girth (HG) and age, body weight (BW), scrotal circumference (SC), testes volume (TV), testis length (TL) and testis width (TW) in bulls maintained at Darabif | 44 |
| 3 | Comparison of heart girth in Kedah-Kelantan bulls maintained at UPM (KK1) and Darabif (KK2) | 45 |
| 4 | Comparison of body weight in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 47 |
| 5 | Correlation and regression between body weight (BW) and age, scrotal circumference (SC), testes volume (TV), testis length (TL), and testis width (TW) in bulls maintained at Darabif | 50 |
| 6 | Comparison of body weight in Kedah-Kelantan bulls maintained at UPM (KK1) and Darabif (KK2) | 51 |
| 7 | Comparison of scrotal circumference in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 54 |
| 8 | Correlation and regression between scrotal circumference (SC), age, testes volume (TV), testis length (TL), and testis width (TW) in bulls maintained at Darabif | 57 |
| 9 | Comparison of scrotal circumference in Kedah-Kelantan bulls maintained at UPM (KK1) and Darabif (KK2) | 58 |



| Table | | Page |
|-------|--|------|
| 10 | Comparison of testes volume in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 60 |
| 11 | Comparison of testes volume in Kedah-Kelantan bulls maintained at UPM (KK1) and Darabif (KK2) | 63 |
| 12 | Comparison of right testis length in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 64 |
| 13 | Correlation and regression between age, testes volume (TV), testis length (TL), and testis width (TW) in bulls maintained at Darabif | 67 |
| 14 | Comparison of right testis length in Kedah-Kelantan bulls maintained at UPM (KK1) and Darabif (KK2) | 68 |
| 15 | Comparison of right testis width in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 70 |
| 16 | Comparison of right testis width in Kedah-Kelantan bulls, maintained at UPM (KK1) and Darabif (KK2) | 74 |
| 17 | Comparison of semen volume (SV), sperm concentration (SC), sperm motility (SM) and live spermatozoa (LS) in Kedah-Kelantan bulls maintained at UPM | 75 |
| 18 | Comparison between puberty and sexual maturity in body size, testis size and semen quality in Kedah-Kelantan, Brahman and crossbred bulls | 77 |
| 19 | Mean (\pm SD) changes in plasma testosterone levels before (B) and after (A) intramuscular administration of GnRH in Kedah-Kelantan bulls at different ages | 78 |

LIST OF FIGURES

| Figure | | Page |
|--------|---|------|
| 1 | Relationship between age and heart girth in Kedah-Kelantan, Brahman and crossbred bulls | 43 |
| 2 | Relationship between age and body weight in Kedah-Kelantan, Brahman and crossbred bulls | 49 |
| 3 | Relationship between age and scrotal circumference in Kedah-Kelantan, Brahman and crossbred bulls | 55 |
| 4 | Relationship between age and testes volume in Kedah-Kelantan, Brahman and crossbred bulls | 61 |
| 5 | Relationship between age and testis length in Kedah-Kelantan, Brahman and crossbred bulls | 66 |
| 6 | Relationship between age and testis width in Kedah-Kelantan, Brahman and crossbred bulls | 72 |
| 7 | Mean plasma testosterone levels before and after intramuscular administration of GnRH in Kedah-Kelantan bulls at 6 to 12 months of age | 79 |
| 8 | Mean plasma testosterone levels before and after intramuscular administration of GnRH in Kedah-Kelantan bulls at 13 to 18 months of age | 81 |
| | Mean plasma testosterone levels before and after intramuscular administration of GnRH in Kedah-Kelantan bulls at 19 to 24 months of age | 82 |



| Figure | | Page |
|--------|--|------|
| 10 | Mean plasma testosterone levels before and after intramuscular administration of GnRH in Kedah-Kelantan bulls at 25 to 30 months of age | 84 |
| 11 | Mean plasma testosterone levels before and after intramuscular administration of GnRH in Kedah-Kelantan bulls at 31 to 36 months of age | 85 |
| 12 | Mean plasma testosterone levels before and after intramuscular administration of GnRH in Kedah-Kelantan bulls at over 36 months of age | 86 |
| 13 | Relationships between age and plasma testosterone levels before and after GnRH treatment, scrotal circumference, testis length and sperm concentration in Kedah-Kelantan bulls | 88 |



LIST OF APPENDICES

| Appendix | | Page |
|----------|---|------|
| 1 | Chemical composition of commercial cattle concentrate | 119 |
| 2 | Cross-reaction of various steroids relative to testosterone | 120 |
| 3 | Mean (\pm SD) age for each period in Kedah-Kelantan, Brahman and crossbred bulls maintained at Darabif | 121 |
| 4 | Mean (\pm SD) Age for each period in Kedah-Kelantan bulls maintained at UPM (KK1) and Darabif (KK2) | 122 |
| 5 | Age changes in body size, testis size, sperm output and plasma testosterone concentration in Kedah-Kelantan bulls maintained at UPM | 123 |
| 6 | Correlations between age, body size, sperm output and plasma testosterone concentration in Kedah-Kelantan bulls maintained at UPM | 124 |
| 7 | Coefficient of variation in heart girth, body weight and scrotal circumference for three breeds of bulls maintained at Darabif | 125 |
| 8 | Ratio between testes volume and heart girth in three breeds of bulls maintained at Darabif | 126 |
| 9 | Ratio between testes volume and body weight in three breeds of bulls maintained at Darabif | 127 |
| 10 | Ratio between testis length and scrotal circumference in three breeds of bulls maintained at Darabif | 128 |



| Appendix | | Page |
|----------|---|------|
| 11 | Ratio between testis width and scrotal circumference in three breeds of bulls maintained at Darabif | 129 |
| 12 | Analysis of variance on effects of breed on heart girth | 130 |
| 13 | Analysis of variance on effects of breed on body weight | 131 |
| 14 | Analysis of variance on effects of breed on scrotal circumference | 132 |
| 15 | Analysis of variance on effects of breed on testes volume | 133 |
| 16 | Analysis of variance on effects of breed on testis length | 134 |
| 17 | Analysis of variance on effects of breed on testis width | 135 |
| 18 | Analysis of variance on effects of breed on semen quality | 136 |
| 19 | Analysis of variance on effects of breed and age on scrotal circumference | 137 |
| 20 | Analysis of variance on effects of breed and age on testes volume | 137 |
| 21 | Analysis of variance on effects of breed and age on testis length | 138 |
| 22 | Analysis of variance on effects of breed and age and testis width | 138 |



LIST OF ABBREVIATIONS

| | | |
|------|---|---------------------------------|
| CV | : | Coefficient of variation |
| ° | : | Degrees centigrade |
| C | : | Degrees centigrade |
| cpm | : | Counter per minute |
| dpm | : | Disintegration per minute |
| FSH | : | Follicle stimulating hormone |
| gm | : | Gram |
| GnRH | : | Gonadotrophin releasing hormone |
| kg | : | Kilogram |
| LH | : | Luteinizing hormone |
| ml | : | Millilitre |
| MW | : | Molecule weight |
| ng | : | Nanogram |
| PGB | : | Phosphate Gelatin Buffer |
| pg | : | Picogram |
| pH | : | Potential hydrogen |
| RIA | : | Radioimmunoassay |
| rpm | : | Rate per minute |
| ug | : | Micogram |
| ul | : | Microlitre |



Abstract of thesis presented to the Senate of Universiti
Pertanian Malaysia in partial fulfilment of the requirements
for the degree of Master of Science

STUDIES IN KEDAH-KELANTAN, BRAHMAN AND
CROSSBRED BULLS AT PUBERTY AND SEXUAL MATURITY

by

ISMAYA

May, 1987

Supervisor : Assoc. Prof. Tan Hock Seng, Ph. D.

Co-Supervisor: Assoc. Prof. Tuan Ariffen Bongso, Ph. D.

Faculty : Veterinary Medicine and Animal Science

Physiological differences within and between breeds could be important in genetic selection and in the planning of management programmes for maximization of reproductive efficiency.

Data for this study were obtained from 21 Kedah-Kelantan bulls (6 to 36 months of age) maintained at Universiti Pertanian Malaysia. Data were also obtained from 109 Kedah-Kelantan bulls and 105 Brahman bulls (6 to 72 months of age) and 25 crossbred bulls (13 to 36 months of age) maintained at Darabif Sdn. Berhad. Heart girth, body weight, scrotal circumference, testes volume, testis length and testis width in all three breeds were measured. Semen production and plasma



testosterone concentration were also investigated in Kedah-Kelantan bulls.

Relationship between age, body size and testis size were found to be significantly different for all three breeds. Brahman bulls were superior to Kedah-Kelantan bulls with crossbred bulls in between. In all three breeds, heart girth increased rapidly between 14 and 21 months of age, besides body weight, scrotal circumference, testes volume, testis length and testis width. At 6 to 12, 13 to 18 and 19 to 24 months of age, 0 per cent, 26.7 per cent and 83.3 per cent of Kedah-Kelantan bulls responded to electroejaculation, respectively. Whereas at over 30 months of age all Kedah-Kelantan bulls responded to electroejaculation. Semen quality was observed to increase with age.

Kedah-Kelantan bulls reached puberty at 482 ± 45 days of age, weighed 151.5 ± 30 kg with a heart girth of 129 ± 8 cm. At puberty, Kedah-Kelantan bulls had a semen volume of 2.6 ± 0.5 ml, sperm concentration of 320 ± 136 millions spermatozoa/ml, sperm motility of 47.5 ± 22.2 per cent and live sperm of 78.3 ± 2.7 per cent. Scrotal circumference, testes volume, testis length and testis width were 21.0 ± 2.6 cm, 282 ± 44 ml, 11.5 ± 1.5 cm and 4.5 ± 0.5 cm, respectively.

At sexual maturity, Kedah-Kelantan bulls had semen volume, sperm concentration, sperm motility and live sperm of 231 per

cent, 218 per cent, 71 per cent and 5 per cent higher than at puberty, respectively. Body weight, heart girth and scrotal circumference in mature bulls were 105 per cent, 33 per cent and 33 per cent higher than at puberty, respectively. Similarly, testes volume, testis length and testis width were 136 per cent, 50 per cent and 37 per cent higher than at puberty, respectively.

Mean testosterone concentration and mean testosterone peak in Kedah-Kelantan bulls increased ($P < 0.01$) with age and fluctuated during 10 hour periods. Mean testosterone concentration and testosterone peak increased from 0.42 to 6.37 ng/ml and 1.27 to 13.70 ng/ml between 6 and 36 months of age, respectively.

Abstrak tesis yang dikemukakan kepada Senat Universiti
Pertanian Malaysia sebagai memenuhi sebahagian daripada syarat
untuk ijazah Master Sains

KAJIAN KEBALIGHAN DAN KEDEWASAAN PADA LEMBU-LEMBU
JANTAN KEDAH-KELANTAN, BRAHMAN DAN KACUKAN

oleh

ISMAYA

Mei, 1987

Penyelia : Prof. Madya Tan Hock Seng, Ph. D.

Penyelia Bersama: Prof. Madya Tuan Ariffee Bongso, Ph. D.

Fakulti : Kedokteran Veterinar dan Sains Peternakan

Perbezaan fisiologi dalam dan antara baka ternakan
memainkan peranan penting dalam pemilihan genetik dan merancang
program pengurusan untuk meningkatkan keberkesanan pembiakan.

Bahan untuk kajian ini diperolehi dari 21 ekor lembu
jantan Kedah-Kelantan yang berumur antara 6 hingga 36 bulan
yang diperlihara di Universiti Pertanian Malaysia, 109 ekor
lembu jantan Kedah-Kelantan, 105 ekor lembu jantan Brahman yang
berumur antara 6 hingga 72 bulan dan 25 ekor lembu kacukan yang
berumur antara 13 hingga 36 bulan yang dipelihara di Darabif
Sdn. Berhad, Pahang. Lilitan dada, berat badan, panjang
keliling kerandut, isipadu buah pelir, panjang dan lebar buah



pelir ketiga-tiga baka telah diukur. Pengeluaran semen dan kadar plasma testosteron lembu Kedah-Kelantan juga telah diselidiki.

Adalah didapati bahawa perhubungan di antara umur, saiz badan dan saiz buah pelir bagi ketiga-tiga jenis baka tersebut sangatlah berbeza. Lembu jantan Brahman lebih unggul daripada jenis Kedah-Kelantan, manakala mutu jenis kacukan pula adalah di antara keduanya. Bagi ketiga-tiga jenis, lilitan dada pesat berkembang di antara umur 14 hingga 21 bulan. Pada waktu yang sama, berat badan, panjang keliling kerandut, isipadu buah pelir serta panjang dan lebarnya juga pesat bertambah. Pada umur 6 hingga 12, 13 hingga 18 dan 19 hingga 24 bulan kadar tindakbalas lembu jantan Kedah-Kelantan terhadap 'electro-ejaculation' ialah 0 peratus, 26.7 peratus dan 83.3 peratus. Pada umur lebih daripada 30 bulan kesemua lembu jantan Kedah-Kelantan membuat tindakbalas terhadap 'electroejaculation'.

Lembu jantan Kedah-Kelantan mencapai tahap baligh pada umur 482 ± 45 hari dengan berat sebanyak 151.5 ± 30 kg serta lilitan dada 129 ± 8 cm. Ketika baligh, lembu Kedah-Kelantan telah menghasilkan semen sebanyak 2.6 ± 0.5 ml, kepekatan spermatozoa 320 ± 136 juta spermatozoa/ml, motiliti sperma 47.5 ± 22.2 peratus dan sperma hidup 78.3 ± 2.7 peratus. Ukuran panjang keliling kerandut, isipadu buah pelir, panjang dan lebar testes ialah 21.0 ± 2.6 cm, 282 ± 44 ml, 11.5 ± 1.5 cm dan 4.5 ± 0.5 cm.

Ketika mencapai kematangan, lembu Kedah-Kelantan mempunyai isipadu semen, kepekatan sperma, motiliti sperma dan sperma hidup sebanyak 231 peratus, 218 peratus, 71 peratus dan 5 peratus lebih tinggi daripada semasa baligh. Berat badan, lilitan dada dan panjang kerandut lembu yang telah mencapai kematangan adalah 105 peratus, 33 peratus dan 33 peratus lebih tinggi daripada ketika baligh. Dalam pada itu, isipadu buah pelir, panjang dan lebar testes adalah 136 peratus, 50 peratus dan 37 peratus lebih tinggi daripada ketika baligh.

Purata kadar testosteron dan purata kemuncak testosteron lembu jantan Kedah-Kelantan bertambah ($P < 0.01$) mengikut umur dan mengalami turun-naik dalam jangkamasa 10 jam. Purata kadar testosteron dan purata kemuncak testosteron bertambah dari 0.42 ke 6.37 ng/ml dan 1.27 ke 13.70 ng/ml pada umur 6 hingga 36 bulan.

CHAPTER I

INTRODUCTION

Kedah-Kelantan cattle, indigenous to Malaysia, represent a potential economic resource for beef cattle production in Malaysia. The animal has gone through a long process of adaptation in Malaysia with high fertility and they constitute a body size which is manageable by local farmers. Moreover, these cattle represent about 82 per cent of the prevailing total cattle population (Devendra and Lee, 1979).

Kedah-Kelantan cattle are relatively small in size with a lower average daily gain compared to Brahman and Brahman cross cattle. Clayton (1983) reported that average daily live weight gain for male animals were 0.36, 0.50 and 0.44 kg/day for Kedah-Kelantan, Brahman and Brahman cross cattle, respectively. The results indicated that Brahman bull could be used to upgrade the potential of Kedah-Kelantan cattle. The approach of crossbreeding will strengthen future breeding strategies to increase beef cattle number and quality.

Puberty may be defined as the point of development when a bull is capable of participating in reproduction. In the present study, puberty was defined as the time when a bull could respond by electroejaculation. Puberty has been defined

as the time when the bull's sexual organs are functionally developed, sexual instincts are prominent and reproduction is possible. Puberty should not be confused with sexual maturity, as the latter is attained at a later date when all systems are functioning at their maximum. Puberty is the time of first breeding potential, while sexual maturity is the time of maximum breeding potential. Pubertal characteristics in Bos indicus bulls have been described by Iqboeli and Rakha (1971) and more recently by Oyedipe et al. (1981), Fields et al. (1982) and Wildeus et al. (1984a). These studies have highlighted the delayed onset of puberty and slower rate of sexual maturation compared to Bos taurus bulls (Fields et al., 1982; Wildeus et al., 1984b). Studies in a tropical environment have shown differences in the development of sperm reserves in Brahman and Sahiwal derived crossbred bulls (Wildeus and Entwistle, 1983).

Furthermore, the lower pregnancy rates achieved with sires of Bos indicus breeding have been attributed to a poorer quality ejaculate (Iqboeli and Rakha, 1971; Fields et al., 1979) and smaller testes (Fields et al., 1979; Blockey, 1980). Besides that, breed groups with average testosterone concentration reached puberty earlier than breed groups of bulls with low testosterone concentration (Lunstra et al., 1978). Testosterone concentration in the first year of life has been studied for beef (Lunstra et al., 1978; Singal and Gomes, 1978) and dairy breeds (Thibier, 1975; Schanbacher,